


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all Aspects of Natural Products Research



Volume 13. Issue 4. Pages 387-512. 2018
ISSN 1934-578X (printed); ISSN 1555-9475
(online) www.naturalproduct.us

EDITOR-IN-CHIEF

DR. PAWAN K AGRAWAL

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS

PROFESSOR MAURIZIO BRUNO

Department STEBICEF,
University of Palermo, Viale delle Scienze,
Parco d'Orleans II - 90128 Palermo, Italy
maurizio.bruno@unipa.it

PROFESSOR CARMEN MARTIN-CORDERO

Department of Pharmacology, Faculty of Pharmacy,
University of Seville, Seville, Spain
carmenmc@us.es

PROFESSOR VLADIMIR I. KALININ

G.B. Elyakov Pacific Institute of Bioorganic Chemistry,
Far Eastern Branch, Russian Academy of Sciences,
Pr. 100-letiya Vladivostoka 159, 690022,
Vladivostok, Russian Federation
kalinin@piboc.dvo.ru

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimaki@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry, University of Wollongong,
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry, Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry, The University of Alabama in Huntsville,
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu

PROFESSOR PING-JYUN SUNG

National Museum of Marine Biology and Aquarium
Checheng, Pingtung 944
Taiwan
pjsung@nmmba.gov.tw

PROFESSOR YASUHIRO TEZUKA

Faculty of Pharmaceutical Sciences, Hokuriku University,
Ho-3 Kanagawa-machi, Kanazawa 920-1181, Japan
y-tezuka@hokuriku-u.ac.jp

PROFESSOR DAVID E. THURSTON

Institute of Pharmaceutical Science
Faculty of Life Sciences & Medicine
King's College London, Britannia House
7 Trinity Street, London SE1 1DB, UK
david.thurston@kcl.ac.uk

HONORARY EDITOR

PROFESSOR GERALD BLUNDEN

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Giovanni Appendino
Novara, Italy

Prof. Norbert Arnold
Halle, Germany

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Vassaya Bankova
Sofia, Bulgaria

Prof. Roberto G. S. Berlinck
São Carlos, Brazil

Prof. Anna R. Bilia
Florence, Italy

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Fatih Demirci
Eskişehir, Turkey

Prof. Francesco Epifano
Chieti Scalo, Italy

Prof. Ana Cristina Figueiredo
Lisbon, Portugal

Prof. Cristina Gracia-Viguera
Murcia, Spain

Dr. Christopher Gray
Saint John, NB, Canada

Prof. Dominique Guillaume
Reims, France

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. Hisaaki Hagiwara
Niigata, Japan

Prof. Judith Hohmann
Szeged, Hungary

Prof. Tsukasa Iwashina
Tsukuba, Japan

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Phan Van Kiem
Hanoi, Vietnam

Prof. Niel A. Koorbanally
Durban, South Africa

Prof. Chiaki Kuroda
Tokyo, Japan

Prof. Hartmut Laatsch
Göttingen, Germany

Prof. Marie Lacaille-Dubois
Dijon, France

Prof. Shoen-Sheng Lee
Taipei, Taiwan

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Stefano Serra
Milano, Italy

Dr. Bikram Singh
Palampur, India

Prof. Marina Stefova
Skopje, Republic of Macedonia

Prof. Leandros A. Skaltsounis
Zografou, Greece

Prof. John L. Sorensen
Manitoba, Canada

Prof. Johannes van Staden
Scottsville, South Africa

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Sylvia Urban
Melbourne, Australia

Prof. Karen Valant-Vetschera
Vienna, Austria

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2017 subscription price: US\$2,595 (Print, ISSN# 1934-578X); US\$2,595 (Web edition, ISSN# 1555-9475); US\$2,995 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Altered Antimicrobial and Anti-biofilm Forming Effect of Thyme Essential Oil due to Changes in Composition

Csilla Gömöri^a, Anita Vidács^c, Erika Beáta Kerekes^a, Elvira Nacs-Farkas^a, Andrea Böszörményi^b, Csaba Vágvolgyi^a and Judit Krisch^c

^aDepartment of Microbiology, Faculty of Science and Informatics, University of Szeged, Közép fasor 52, Szeged, H-6726, Hungary

^bSemmelweis University, Department of Pharmacognosy, Üllői út 26, Budapest, H-1085, Hungary

^cInstitute of Food Engineering, Faculty of Engineering, University of Szeged, Mars tér 7, Szeged, H-6724, Hungary

gmri.csilla@gmail.com

Received: October 6th, 2017; Accepted: February 25th, 2018

The antimicrobial and anti-biofilm forming effect of thyme (*Thymus vulgaris*) essential oil (TEO) with different compositions was evaluated. Normally the main component in this TEO is thymol, but in 2014 we found that the proportions of γ -terpinene and *p*-cymene (the precursors in thymol biosynthesis) increased and that of thymol decreased. This altered composition led to changes in the antimicrobial and anti-biofilm forming capacity of the essential oil depending also on the type of microorganisms. In the case of bacteria, minimal inhibitory (MIC) and minimal bactericidal (MBC) concentrations either decreased or increased. In the case of yeasts, minimal fungicidal concentrations (MFC) increased 4- and 8-fold for TEO containing *p*-cymene as the main component. On the contrary, MIC values decreased for all the tested moulds. Anti-biofilm forming activity of TEO containing *p*-cymene as its main component decreased in almost all cases and *P. fluorescens* biofilm forming capacity was even enhanced.

Keywords: Antimicrobial effect, Biofilms, Changed composition, Thyme, *Thymus vulgaris*, Essential oils.

The yield and quality of essential oils (EOs) and extracts depend on the edaphic, climatic conditions and the genotype of the plant. Geographical location, altitude, soil properties and irrigation influence the phytochemical composition and antioxidant activity of EOs. The chemical composition of EOs can be highly variable among cultivars, thus it is important to optimize both growing conditions and the genotype [1]. Furthermore, it is necessary to know the optimal harvesting time in order to provide high quality EOs [2].

EO yield is prominently influenced by the phenological stage (vegetative stage, full bloom, seed maturation) of the plant, but plant age [3, 4], leaf age and leaf positions [5] are also important. Seasonal changes [6], growth stages, origin of the herb [7], and the method of extraction can also influence yield, product quality, and quantitative and qualitative characteristics of EOs [8].

Usano-Aleman *et al.* studied EO yields and qualities of *Salvia lavandulifolia* and confirmed that plants in the vegetative and flowering stages have higher content of terpene hydrocarbons than those at the seed maturation stage, when the content of oxygenated derivatives increased remarkably [3].

The main components of *Thymus* EOs are thymol, *p*-cymene, γ -terpinene or carvacrol, and linalool, depending on the chemotype of the oil [9, 10, 11, 12].

The chemical compositions of EOs of *Thymus* species are also affected by environmental conditions (e.g., humidity, temperature, radiation, soil characteristics, moisture availability, fertility, electrical conductivity) and agronomic management practices. Higher elevation and colder temperature provided better growing conditions, and a higher accumulation of EO in thyme leaves [13].

Unsuitable environmental conditions may limit photosynthesis and change nutrient uptake in plants. Under stress (e.g., bad weather conditions, limited nutrients) they are more susceptible to pests and pathogens. During cold months with few hours of sunlight, decreased EO production was noticed [13].

Results of an experiment conducted in 2010 showed that changes in quality and quantity of EOs are species-specific under the same circumstances. The composition of EOs could change with the water supply. Using *Ocimum basilicum* L. as a test plant, the content of linalool decreased, but that of 1,8-cineol and tau-cadinol increased with increasing water scarcity. In the case of *Satureja hortensis*, the level of γ -terpinene increased, but that of carvacrol decreased with increased water supply [14].

EOs are used in the food industry primarily as flavorings. They also have a potential food preservative ability because they are effective against food spoilage bacteria and mycotoxin producing fungi [15]. EOs could be ideal food additives instead of synthetic ones [16] as they are natural and mostly non-toxic to humans [17]. Most EOs have antibacterial activity against pathogens [18]. Besides this, studies show that they are also effective against bacterial biofilms [19, 20]. Biofilm formation is common among bacteria; in this form, they become more resistant to antimicrobials and sanitizing agents [21]. The antimicrobial activity of EOs is attributed to their components [22], which are generally classified into two groups of dissimilar biosynthetic origin. The main group contains terpenes and terpenoids, whereas the other one consists of aromatic compounds (phenylpropanoids) [23, 24].

The phenolic components are chiefly responsible for the antimicrobial effect as membrane permeabilizers. Gram-positive microorganisms are generally more sensitive to EOs than Gram-negative ones [12, 20, 25].

Table 1: Chemical composition (%) of the thyme essential oils tested.

Compound	Chemical composition (%)			
	2013		2014	
	Retention time (min)	Component proportion (%)	Retention time (min)	Component proportion (%)
1,8-Cineol	8.0-8.1	9.8	-	-
3-Carene	-	-	7.3	0.5
Borneol	11.0	1.0	10.7	0.5
Camphene	6.3	0.9	6.2	4.9
Carvacrol	13.3	3.2	13.2	3.1
Caryophyllene oxide	-	-	17.7	0.2
Isoborneol	-	-	10.6	0.4
Linalool	9.5	5.1	9.3	5.3
Linalyl acetate	-	-	12.4	0.3
Myrcene	-	-	6.9	4.0
Nerol	-	-	12.6	0.1
Neryl acetate	-	-	14.4	0.3
<i>p</i> -Cymene	7.9	22.1	7.8	30.1
Terpinen-4-ol	11.5	0.6	10.9	0.9
Terpinolene	-	-	9.0	0.9
Thymol	13.2	46.3	13.1	23.2
α -Humulene	-	-	15.8	0.2
α -Pinene	5.9	0.9	5.8	3.2
α -Terpinene	7.7	2.2	7.6	5.9
α -Terpineol	-	-	11.4	0.6
β -Caryophyllene	15.4	2.5	15.2	2.5
β -Pinene	6.9	1.4	6.7	0.8
γ -Terpinene	8.6	0.3	8.5	10.0
Terpene hydrocarbons		30.3		62.1
Terpenoids		66.0		34.9
Total		96.3		97.0

Our group has been working with EOs for years. We worked with a thymol chemotype of *Thymus vulgaris* L. EO in which thymol always formed the highest proportion of the main components. In 2014, after a rainy summer, the composition of the TEO changed dramatically; this initiated us to examine if changes in the composition of the essential oil had had an influence on its antimicrobial efficiency.

Table 1 presents the chemical compositions of *T. vulgaris* EO from different years. Chromatographic analysis showed that the major component of thyme EO in 2013 was thymol and in 2014 it was its precursor, *p*-cymene. 1-8 Cineol was detected only in 2013, while the content of γ -terpinene was much higher in the following year.


In most cases, the MIC and MBC values changed in 2014, except for *Bacillus subtilis* for which no changes were observed. MIC and MBC values for *Staphylococcus aureus* and *Pseudomonas fluorescens* increased compared with values in 2013 (Table 2).

Table 2: MIC and MBC values of TEO with original and altered composition against bacteria (mg/mL).

Bacteria	MIC (mg/mL)		MBC (mg/mL)	
	2013	2014	2013	2014
<i>E. coli</i>	1.6	1.6	1.6	32.0
<i>P. putida</i>	20.0	8.0	>100	8.0
<i>P. fluorescens</i>	1.6	2.0	1.6	3.2
<i>B. cereus</i>	1.6	0.8	3.2	1.6
<i>B. subtilis</i>	0.8	0.8	1.6	1.6
<i>S. aureus</i>	0.8	1.6	1.6	3.2
MRSA	3.2	1.6	12.5	3.2
<i>L. monocytogenes</i>	1.6	3.2	6.3	6.3

The MBC of *Escherichia coli* increased twenty times, while that of *Pseudomonas putida* decreased by about one tenth compared with 2013 values. MIC and MBC values of MRSA and *B. cereus* var. *mycoides* decreased moderately.

Sessile cells in biofilms of most investigated bacteria showed different results compared with planktonic cells. Almost in all cases, thyme EO from 2013 was more effective than that from 2014 (Figure 1). For *B. cereus* var. *mycoides* there was no significant difference between the two Thyme EO preparations, but in the case

**Figure 1:** Anti-biofilm forming effect of TEO from 2013 and 2014. The oil was used in MIC/2 concentration. A: *B. cereus*; B: *B. subtilis*; C: *L. monocytogenes*; D: *P. putida*; E: *E. coli*; F: *P. fluorescens*; G: *P. anomala*.

of *E. coli* the EO containing more *p*-cymene (2014) was more effective. *P. fluorescens* biofilms were enhanced by the EO, which is in agreement with the findings of Sandasi et al. where monoterpenes enhanced biofilm formation [26]. It seems that by the accumulation of *p*-cymene and the reduction of thymol the biofilm formation was stimulated instead of inhibited.

In the case of yeasts, the MFC values increased 4-, and 8-fold from 2013 to 2014. On the contrary, MIC values decreased for all the tested moulds (Table 3).

Table 3: MFC values of thyme EO with original and altered composition tested *in vitro* against yeasts and moulds (mg/mL).

Fungi	Year	
	2013	2014
<i>Candida albicans</i>	0.78	3.12
<i>Candida parapsilosis</i>	0.19	1.56
<i>Pichia anomala</i>	1.56	1.56
<i>Aspergillus fumigatus</i>	0.39	<0.19
<i>Aspergillus terreus</i>	0.78	0.39
<i>Fusarium solani</i>	0.78	0.39
<i>Rhizopus microsporus</i>	0.19	<0.19

In all cases, thymol alone showed better antimicrobial activity than *p*-cymene. Mixing the two components resulted in mainly a stepwise increase in the MIC and MBC values (Table 4). For staphylococci, the first increase occurred at the ratio of thymol: *p*-cymene of 4:1, while for the other bacteria the ratio was either 1:1 or 3:2. MIC and MBC values of the compounds either alone or in combination were in the same range as those of the whole EO.

Table 4: MIC/MBC values (mg/mL) after treatment with thymol and *p*-cymene in different proportions.

Thymol: <i>p</i> -Cymene (v:v)	<i>E. coli</i>		<i>S. aureus</i>		MRSA		<i>L. monocytogenes</i>		<i>B. subtilis</i>		<i>B. cereus</i>	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
100:0	0.20	0.40	0.10	0.20	0.10	0.20	0.40	0.80	<0.05	0.05	0.40	0.80
80:20	0.20	0.40	0.20	0.40	0.20	0.40	0.40	0.80	<0.05	0.05	0.40	0.80
60:40	0.20	0.80	0.20	0.40	0.20	0.40	0.40	0.80	0.05	0.10	0.40	0.80
50:50	0.20	0.80	0.20	0.40	0.20	0.40	0.80	1.60	0.20	0.40	0.80	0.80
40:60	0.40	0.80	0.20	0.40	0.20	0.40	0.80	1.60	0.20	0.40	0.80	1.60
20:80	0.80	1.60	0.20	0.40	0.40	0.80	1.60	3.20	0.20	0.40	0.80	1.60
0:100	1.60	3.20	0.80	1.60	0.80	1.60	3.20	>25.00	0.80	1.60	0.80	1.60

Several authors reported the importance of different environmental factors to the EO compositions of plants [2, 14, 27]. Drought stress, but also too much rain and little sunshine, can lead to changes in the biochemical pathways of the synthesis of EO compounds. In Hungary (and in the whole of Central Europe), summer showed a lot of extremities in 2013 and 2014. Temperature increased from 2013 to 2014, but in contrast to the global trend, in Hungary 2014 was rainier than 2013.

It seems that this unusual climate of 2014 affected the EO biosynthesis of *Thymus* plants. Thymol is synthesized from *p*-cymene, which in turn is derived from γ -terpinene [28]. In the EO from 2014 the precursors: *p*-cymene and γ -terpinene were found in large amounts while thymol was reduced to about a half (Table 1).

This altered composition led to changes in the antimicrobial and anti-biofilm forming activity of the oil. The data for the antimicrobial effect obtained with *p*-cymene and thymol alone and in combinations (Table 4) showed that thymol alone had better antimicrobial activity than *p*-cymene either alone or in any combination of the two. Dorman and Deans also described the relative inefficiency of *p*-cymene compared with that of thymol [22]. However, the EO with decreased thymol and increased *p*-cymene and γ -terpinene content (year 2014) showed, in certain cases, higher antimicrobial activity than the EO from 2013. This indicated that not only the main ingredients had a crucial role in the antimicrobial activity, but the individual sensitivity of the used strains was also very important.

In 2014, the proportion of terpenes increased at the expense of terpenoids. It is generally assumed that terpenoids have better antimicrobial effect than terpenes [22, 23], but synergistic interactions among monoterpene hydrocarbons and terpenoids cannot be excluded.

This study investigated the antimicrobial and anti-biofilm forming effects of thyme essential oils from plants harvested in two consecutive years. The major component of the TEO in 2013 was thymol and in 2014 (with hot and rainy weather), its precursor, *p*-cymene. The altered composition led to remarkable changes in the antimicrobial and anti-biofilm forming capacity of TEO. Thymol alone had better antimicrobial effect than *p*-cymene alone or in any combination of the two, but the minor components in the whole TEO could modulate this effect.

An important conclusion of our finding is that, as plant EO compositions cannot be standardized, MIC values need to be established for every new batch of EO.

Experimental

Test organisms: Selected bacteria and fungi from the Szeged Microbiology Collection (SZMC; WDCM987) and American Type Culture Collection (ATCC; WDCM1) were investigated. The investigated bacteria were: *Bacillus cereus* var. *mycoides* SZMC 0042, *B. subtilis* SZMC 0209, *Escherichia coli* SZMC 0582,

Listeria monocytogenes SZMC 21307, *Pseudomonas putida* SZMC 291T, *P. fluorescens* SZMC 16106, *Staphylococcus aureus* ATCC 43300 (methicillin resistant, (MRSA) and ATCC 25923 (antibiotic sensitive).

S. aureus and *E. coli* strains were grown in LB broth: 10 g NaCl (VWR, Belgium), 10 g bacto peptone (Oxoid, England), 5 g yeast extract (HiMedia, India) in 1000 mL. *B. cereus*, *B. subtilis*, and *P. putida* were cultivated in TGE broth: 10 g glucose (VWR, Belgium), 5 g bacto peptone, 2.5 g yeast extract, in 1000 mL. TSB was used for *L. monocytogenes*: 17 g casein peptone (Merck, Germany), 3 g soy peptone (Oxoid, England), 2.5 g glucose, 5 g NaCl, 2.5 g K₂HPO₄ (Reanal, Hungary), in 1000 mL. *E. coli*, *S. aureus*, and *L. monocytogenes* were incubated at 37°C, *P. putida* at 25°C, and bacilli at 30°C for 18-20 h to reach early stationary growth phase.

The tested fungal species were: *Aspergillus fumigatus* SZMC 2394, *A. terreus* SZMC 2414, *Fusarium solani* SZMC 11412, *Rhizopus microsporus* SZMC 13644, *Candida albicans* SZMC 1363, *C. parapsilosis* SZMC 1408, and *Pichia anomala* SZMC 8061. For the cultivation of yeasts and moulds malt extract agar was used (0.4% malt extract, 0.1% yeast, 1% glucose, and 2% agar).

Essential oils and components: Thyme (*Thymus vulgaris*) essential oils produced in 2013 and 2014 were purchased from Aromax Natural Products Ltd. (Budapest, Hungary). EOs were prepared by steam distillation of aerial parts of the plants.

Thymol and *p*-cymene were purchased from Sigma Aldrich (Budapest, Hungary).

Identification of essential oil compounds

Solid phase microextraction (SPME) conditions: Essential oil (0.5 mL) was put into vials (20 mL headspace) and sealed with a silicon/PTFE septum prior to SPME-GC/MS analysis. Sample preparation using a static headspace solid phase microextraction (sHS-SPME) technique was carried out with a CTC Combi PAL (CTC Analytics AG, Zwingen, Switzerland) automatic multipurpose sampler using a 65 μ M StableFlex polydimethylsiloxane/divinyl benzene (PDMS/DVB) SPME fiber (Supelco, Bellefonte, PA, USA). After an incubation period of 5 min at 100°C, extraction was performed by exposing the fiber to the headspace of a 20 mL vial containing the plant material for 20 min at 100°C. The fiber was then immediately transferred to the injector port of the GC/MS, and desorbed at 250°C for 1 min. The SPME fiber was cleaned and conditioned in a Fibre Bakeout Station in a pure nitrogen atmosphere at 250°C for 15 min.

GC-MS conditions: The analyses were carried out with an Agilent 6890N/5973N GC-MSD (Santa Clara, CA, USA) system equipped with an Agilent SLB-5MS capillary column (30 m \times 250 μ m \times 0.25 μ m). The GC oven temperature was programmed to increase from 60°C (3 min isothermal) – 250°C at 8°C/min (1 min isothermal). High purity helium (6.0) was used as carrier gas at 1.0 mL/min

(37 cm/s) in constant flow mode. The injector temperature was 250°C and the split ratio was 1:50. The mass selective detector was equipped with a quadrupole mass analyzer and was operated in electron ionization mode at 70 eV in full scan mode (41–500 amu at 3.2 scan/s). The data were evaluated using MSD ChemStation D.02.00.275 software (Agilent). The identification of the compounds was carried out by comparing retention times and recorded spectra with the data of authentic standards, and the NIST 2.0 library was also consulted.

Determination of minimum inhibitory and minimum bactericide concentrations: For minimum inhibitory concentration (MIC) determination, the broth microdilution method was used. In 96-well microtiter plates (Costar, USA) 100 µL cell suspensions (10^5 CFU/ml) were mixed with 100 µL EO solution containing 0.1% TWEEN 40 in the concentration range between 0.1–100 mg/mL. Negative controls were EOs in sterile medium with TWEEN 40, and positive controls were inoculated media without EO. After 18–24 h incubation at the appropriate temperatures, growth of the bacteria was determined after staining with 20 µL resazurin (Cell Titer-Blue® Reagent, Promega) indicator. Color was checked after 1 h. Oxido-reductase enzymes of living cells reduce the blue colored resazurin to the pink colored resorufin [29]. MIC was determined as the lowest concentration where the blue colour did not change. Six replicates were prepared for each EO concentration.

Minimum bactericide concentration (MBC) was detected by the tracking plate method [30] transferring 10 µL from the microplate wells corresponding to MIC and higher concentrations on suitable medium. After 24 h incubation at appropriate temperature, the appearance of bacterial colonies was checked. MBC was defined as the EO concentration where no colony growth was observed.

Biofilm formation and treatment: Microtiter plates (96-well) were inoculated with 200 µL of 24-h-old liquid culture (10^8 cfu/mL). Following 4 h of cell adhesion, the supernatant was removed, and the plates were rinsed with physiological saline. Fresh medium (200 µL) containing TEO was added in MIC/2 concentration to each well, and the plates were further incubated for 24 h. Positive controls contained the inoculated growth medium without any TEO, and negative controls contained TEO in growth medium. Biofilms were visualized after crystal violet staining, as described by Kerekes *et al.* [19]. Experiments were repeated at least 2 times with 6 parallel measurements.

Determination of minimal fungicidal concentration: Minimal fungicidal concentration (MFC) values were determined by macrodilution for yeasts, and by food poisoning method for moulds, as described in the literature [31].

For yeasts, test tubes contained 1 mL malt extract medium and 1 mL EO solution to obtain final concentrations of 25 - 0.2 mg/mL in a twofold dilution range. Tween 40 (1%) was used for the solubilization of EOs. Test tubes were inoculated with 0.1 mL of 24-h-old yeast suspension (10^5 cfu/mL). After 24-h incubation, the growth was checked by the tracking plate method, as described earlier. Plates were incubated at 30°C for 24 h, and then the number of colonies was evaluated. MFC was defined as the EO concentration where no colony growth was observed.

Spore suspensions (10 µL, 10^5 cfu/mL) from 72-h-old mould cultures were dropped onto plates containing 15 mL medium and 0.1% Tween-40 supplemented with EOs in the concentration range of 25–0.2 mg/mL. Petri dishes were sealed with parafilm and incubated at 25°C for 7 days. Control plates (without EOs) were inoculated by the same procedure. After 7 days of incubation the MFC was defined as the minimum EO concentration at which no visible growth was observed. For each concentration 3 parallel measurements were carried out.

Antimicrobial effect and interaction of thymol and p-cymene: The antimicrobial effect of thymol and its precursor p-cymene was investigated alone and in combinations. The tested ratios were (thymol: p-cymene): 4:1, 3:2, 1:1, 2:3, and 1:4. Stock solutions (100 mg/mL) were prepared in 96% ethanol. Antimicrobial effect was determined by the broth microdilution method. Wells containing the components alone or in combination mixed with the appropriate media were inoculated with bacteria (10^5 cfu/mL) and incubated for 24 h. After that, absorbance was measured at 595 nm (SPECTROstar microplate reader, BMG Labtech). The concentration at which absorbance decreased below 10% of the positive control was considered as MIC.

Statistical analysis: Data were analyzed using one-way ANOVA (GraphPad Prism 6.0). Means were compared by the Tukey's test and level of significance was $p < 0.05$.

Acknowledgments - This work was connected to the projects GINOP-2.3.2-15-2016-00012 (Széchenyi 2020 Programme) and TÉT 10-1-2013-0019. Elvira Nacs-Farkas thanks project NTP-NFTÖ-16-450.

References

- [1] Formisano C, Delfino S, Oliviero F, Tenore GC, Rigano D, Senatore F. (2015) Correlation among environmental factors. Chemical composition and antioxidative properties of essential oil and extracts of chamomile (*Matricaria chamomilla* L.) collected in Molise (South-central Italy). *Industrial Crops and Products*, **63**, 256–263.
- [2] Kakaraparthi PS, Srinivas KVNS, Kumar JK, Kumar AN, Rajput DK, Sarma VUM. (2014) Variation in the essential oil content and composition of *Citronella* (*Cymbopogon winterianus* Jowitt) in relation to time of harvest and weather conditions. *Industrial Crops and Products*, **61**, 240–248.
- [3] Usano-Aleman J, Palá-Paúl J, Herraiz-Peñalver D. (2016) Essential oil yields and qualities of different clonal lines of *Salvia lavandulifolia* monitored in Spain over four years of cultivation. *Industrial Crops and Products*, **80**, 251–261.
- [4] Tahmasebi S, Majd A, Mehrafarin A, Jonoubi P. (2016) Comparative ontogenetic survey of the essential oil composition in *Origanum vulgare* L., and *Origanum majorana* L. *Acta Biologica Szegediensis*, **60**, 105–111.
- [5] Gupta KA, Ganjewala D. (2015) A study on developmental changes in essential oil content and composition in *Cymbopogon flexuosus* cultivar *Suvarna*. *Acta Biologica Szegediensis*, **59**, 119–125.
- [6] Wang H, Liu Y, Wei S, Yan Z. (2012) Comparative seasonal variation and chemical composition of essential oils from the leaves and stems of *Schefflera heptaphylla* using microwave-assisted and conventional hydrodistillation. *Industrial Crops and Products*, **36**, 229–237.
- [7] Sellami IH, Maamouri E, Chahed T, Wannes WA, Kchouk ME, Marzouk B. (2009) Effect of growth stage on the content and composition of the essential oil and phenolic fraction of sweet marjoram (*Origanum majorana* L.). *Industrial Crops and Products*, **30**, 395–402.
- [8] Memarzadeh SM., Pirbalouti AG, AdibNejad M. (2015) Chemical composition and yield of essential oils from Bakhtiari savory (*Satureja bachtiarica* Bunge.) under different extraction methods. *Industrial Crops and Products*, **76**, 809–816.
- [9] Jia HL, Ji QL, Xing SL, Zhang PH, Zhu GL, Wang XH. (2010) Chemical composition and antioxidant, antimicrobial activities of the essential oils of *Thymus marschallianus* Will. and *Thymus proximus* Serg. *Journal of Food Science*, **75**, 59–65.

- [10] Iten F, Saller R, Abel G, Reichling J. (2009) Additive antimicrobial effects of the active components of the essential oil of *Thymus vulgaris* – Chemotype carvacrol. *Planta Medica*, **75**, 1231-1236.
- [11] Abdelli W, Bahri F, Romane A, Höferl M, Wanner J, Schmidt E, Jirovetz L. (2017) Chemical composition and anti-inflammatory activity of Algerian *Thymus vulgaris* essential oil. *Natural Product Communications*, **12**, 611-614.
- [12] Schmidt E, Wanner J, Hiifer M, Jirovetz L, Buchbauer G, Gochev V, Girova T, Stoyanova A, Geissler M. (2012) Chemical composition, olfactory analysis and antibacterial activity of *Thymus vulgaris* chemotypes geraniol, 4-thujanol/terpinen-4-ol, thymol and linalool cultivated in southern France. *Natural Product Communications*, **7**, 1095-1098.
- [13] Pirbalouti AG, Hashemi M, Ghahfarokhi FT. (2013) Essential oil and chemical compositions of wild and cultivated *Thymus daenensis* Celak and *Thymus vulgaris* L. *Industrial Crops and Products*, **48**, 43-48.
- [14] Radácsi P, Inotai K, Sárosi SZ, Czövek P, Bernáth J, Németh É. (2014) Effect of water supply on the physiological characteristic and production of basil (*Ocimum basilicum* L.). *European Journal of Horticultural Science*, **75**, 193-197.
- [15] Nguefack J, Leth V, Amvam Zollo PH, and Mathur SB. (2004) Evaluation of five essential oils from aromatic plants of Cameroon for controlling food spoilage and mycotoxin producing fungi. *International Journal of Food Microbiology*, **94**, 329-334.
- [16] Shukla R, Singh P, Prakash B, Dubey NK. (2012) Efficacy of *Acorus calamus* L. essential oil as a safe plant-based antioxidant, Aflatoxin B₁ suppressor and broad spectrum antimicrobial against food-infesting fungi. *International Journal of Food Science & Technology*, **48**, 128-135.
- [17] Soliman KM, Badeaa RI. (2002) Effect of oil extracted from some medicinal plants on different mycotoxigenic fungi. *Food and Chemical Toxicology*, **40**, 1669-1675.
- [18] Lambert RJ, Skandamis PN, Coote PJ, Nychas GJ. (2001) A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *Journal of Applied Microbiology*, **91**, 453-462.
- [19] Kerekes EB, Deák É, Takó M, Tserennadmid R, Petkovits T, Vágvolgyi Cs, Krisch J. (2013) Anti-biofilm forming and anti-quorum sensing activity of selected essential oils and their main components on food-related micro-organisms. *Journal of Applied Microbiology*, **115**, 933-942.
- [20] Burt SA. (2004) Essential oils: their antibacterial properties and potential applications in foods: a review. *International Journal of Food Microbiology*, **94**, 223-253.
- [21] Prakash B, Veeregowda BM, Krishnappa G. (2003) Biofilms: a survival strategy of bacteria. *Current Science*, **85**, 1299-1307.
- [22] Dorman HJD, Deans SG. (2000) Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *Journal of Applied Microbiology*, **88**, 308-316.
- [23] Bakkali F, Averbeck S, Averbeck D, Idaomar M. (2008) Biological effects of essential oils - a review. *Food and Chemical Toxicology*, **46**, 446-475.
- [24] Jayasena DD, Jo C. (2013) Essential oils as potential antimicrobial agents in meat and meat products: A review. *Trends in Food Science & Technology*, **34**, 96-108.
- [25] Demuner AJ, Barbosa LCA, Gonçalves Magalhaes C, da Silva CJ, Alvares Maltha CR, Pinheiro AL. (2011) Seasonal variation in the chemical composition and antimicrobial activity of volatile oils of three species of *Leptospermum* (Myrtaceae) grown in Brazil. *Molecules*, **16**, 1181-1191.
- [26] Sandasi M, Leonard CM, Viljoen AM. (2008) The effect of five common essential oil components on *Listeria monocytogenes* biofilms. *Food Control*, **19**, 1070-1075.
- [27] Llorens-Molina JA, Vacas S. (2016) Effect of drought stress on essential oil composition of *Thymus vulgaris* L. (Chemotype 1, 8-cineole) from wild populations of Eastern Iberian Peninsula. *Journal of Essential Oil Research*, **29**, 145-155.
- [28] Poulose AJ, Croteau R. (1978) Biosynthesis of aromatic monoterpenes: conversion of γ -terpinene to *p*-cymene and thymol in *Thymus vulgaris* L. *Archives of Biochemistry and Biophysics*, **187**, 307-314.
- [29] Sarker SD, Nahar L, Kumarasamy Y. (2007) Microtitre plate-based antibacterial assay incorporating resazurin as an indicator of cell growth, and its application in the *in vitro* antibacterial screening of phytochemicals. *Methods*, **42**, 321-324.
- [30] Jett BD, Hatter KL, Huycke MM, Gilmore MS. (1997) Simplified agar plate method for quantifying viable bacteria. *BioTechniques*, **23**, 648-650.
- [31] Horváth Gy, Török Jenei J, Vágvolgyi Cs, Böszörményi A, Krisch J. (2016) Effects of essential oil combinations on pathogenic yeasts and moulds. *Acta Biologica Hungarica*, **67**, 205-214.

Characteristic Aromas of *Quercus crispula* (Japanese Oak)

Toshio Hasegawa, Ryosuke Mamada, Takahisa Yamanaka, Bujiyuro Shimazaki and Takashi Fujihara 475

Microbial Transformation of *Calamintha glandulosa* Essential Oil by *Aspergillus niger*

Miroslav Novakovic, Danka Bukvicki, Vlatka Vajs, Vele Tesevic, Slobodan Milosavljevic, Petar Marin and Yoshinori Asakawa 479

Altered Antimicrobial and Anti-biofilm Forming Effect of Thyme Essential Oil due to Changes in Composition

Csilla Gömöri, Anita Vidács, Erika Beáta Kerekes, Elvira Nacsá-Farkas, Andrea Böszörményi, Csaba Vágvölgyi and Judit Krisch 483

Accounts/Reviews

Chemo- and Enantioselective Catalytic Hydrogenation of α,β -Unsaturated Ketones and Aldehydes as a Tool to Introduce Chiral Centers at α - or β -Positions of Ketones

Hisahiro Hagiwara 489

A Phytochemical and Pharmacological Review on *Cynomorium songaricum* as Functional and Medicinal Food

Jin-Long Cui, Yi Gong, Xiao-Zan Xue, Yan-Yan Zhang, Meng-Liang Wang and Jun-Hong Wang 501

Natural Product Communications

2018

Volume 13, Number 4

Contents

<u>Original Paper</u>	<u>Page</u>
Spatial Structure and Antimicrobial Activity of Cyclopropane Derivative of Limonene Daniyar Sadyrbekov, Timur Saliev, Yuri Gatilov, Ivan Kulakov, Roza Seidakhmetova, Tulegen Seilkhanov and Sholpan Askarova	387
Chemical and Genetic Identity of <i>Ligularia tsangchanensis</i> and <i>L. muliensis</i>. Isolation of a Cacalol Precursor from a Hybrid of <i>L. tsangchanensis</i> and <i>L. vellerea</i> Chiaki Kuroda, Taketo Itoh, Yurika Suzuki, Yuko Suzuki, Yasuko Okamoto, Motoo Tori, Ryo Hanai and Xun Gong	389
Sesquiterpene Lactones and Phenolics from Roots of <i>Leontodon hispidus</i> subsp. <i>hispidus</i> Klaudia Michalska, Katarzyna Pieron and Anna Stojakowska	393
Synthesis and Insecticidal Activity of β-Dihydroagarofuran Acetal Derivatives Xin Xi, Ximei Zhao, Feng Zhu, Jielu Wei, Zhan Hu, Wenjun Wu and Jiwen Zhang	395
Isolation, and Structural Characterization of a New Hexa β-D-Glucopyranosyl Diterpene from <i>Stevia rebaudiana</i> Venkata Sai Prakash Chaturvedula and Srinivasa Rao Meneni	399
Phytochemistry of <i>Weigela</i> x "<i>kosteriana variegata</i>" (Caprifoliaceae) Nampoina Andriamisaina, Anne-Claire Mitaine-Offier, Benoist Pruvot, Johanna Chluba, Tomofumi Miyamoto, Chiaki Tanaka and Marie-Aleth Lacaille-Dubois	403
New Acetylated Saponins from the Leaves of <i>Trevesia palmata</i> Le Thi Thanh Thao, Do Quyen, Duong Binh Vu, Bui Huu Tai and Phan Van Kiem	407
A New Saponin with Anti-HIV-1 Protease Activity from <i>Acacia pennata</i> Van-Dung Nguyen, Hong-Loan Thi Nguyen, Linh-Chi Do, Vu Van Tuan, Phuong Thien Thuong and Tuan-Nghia Phan	411
Ginsenoside Rg1 Exhibits Anti-asthmatic Activity in an <i>Aspergillus</i> Protease-Induced Asthma Model in Mice Chiyeon Lim, Sehyun Lim, Byoungcho Lee and Suin Cho	415
Daffodils as Potential Crops of Biologically-active Compounds: Assessment of 40 Ornamental Taxa for their Alkaloid Profile and Cholinesterases Inhibition Activity Kateřina Breiterová, Miroslav Ločárek, Eliška Kohelová, Markéta Talácková, Daniela Hulcová, Lubomír Opletal and Lucie Čahlíková	419
Antifouling Potentials and Metabolite Profiles of Two Marine-derived Fungal Isolates Xiao-Yong Zhang, Muhammad Amin, Xin-Ya Xu and Shu-Hua Qi	423
New Flavanol Dimers from the Bark of <i>Celtis tetrandra</i> and Their TRAIL Resistance-Overcoming Activity Prapairot Seephonkai, Ryusho Ishikawa, Midori A. Arai, Thaworn Kowithayakorn, Takashi Koyano and Masami Ishibashi	427
Anti-oxidative Fraction of <i>Lycorma delicatula</i> Alleviates Inflammatory Indicators Seung-hwa Baek, Okjin Joung, Ho-Young Lee, Jae-Cheon Shin, Won-Sik Choi, Tae Hun Lee, Jae-Sam Hwang, Sung-Hee Nam, Hyeon-U Son and Sang-Han Lee	431
A New 3-Prenyl-2-flavene and Other Extractives from <i>Baphia massaiensis</i> and Their Antimicrobial Activities Ngonye Keroletswe, Runner R. T. Majinda and Ishmael B. Masesane	435
<i>In vitro</i> Shoot Cultures of <i>Pterospartum tridentatum</i> as an Alternative to Wild Plants as a Source of Bioactive Compounds José Carlos Gonçalves, Maria Teresa Coelho, Maria da Graça Diogo, Vitor D. Alves, Maria Rosário Bronze, Manuel António Coimbra, Vitor Manuel Martins and Margarida Moldão-Martins	439
Effect of Pretreatment of Morin on the Cisplatin-Induced Toxicity on LLC-PK1 and T24 Cells Claudia Bello-Alvarez, Ángela Patricia Moreno-Londoño and José Pedraza-Chaverri	443
Cancer Chemopreventive Activity of Xanthones from <i>Calophyllum elatum</i> Chihiro Ito, Takuya Matsui, Takashi Kobayashi, Harukuni Tokuda, Sivabalan Shanmugam and Masataka Itoigawa	447
A New Secondary Metabolite from Korean Traditional Herb Plant <i>Hovenia dulcis</i> Tu Cam Le, Kyung-Yun Kang, Inho Yang, Alain S. Leutou, Jaeyoung Ko, Young-Jin Son, Sung-Tae Yee and Sang-Jip Nam	451
Resveratrol Oligosaccharide Induces mRNA Expression for <i>SIRT</i> Hiroki Hamada, Kei Shimoda, Yasukazu Saitoh, Shouta Doi, Yuya Fujitaka, Tsubasa Ono, Hatsuyuki Hamada and Minami Araki	455
Ultrasound-assisted Extraction of Polysaccharide from Radix Bupleuri using Response Surface Methodology and Antioxidant Research of the Polysaccharide Changchun Hao, Hongjin Qu, Shi Chen, Wenjing Han and Runguang Sun	457
α-Glucosidase Inhibitor Isolated from <i>Blechnum pyramidatum</i> Lilia Cherigo and Sergio Martínez-Luis	461
<i>Cordyceps militaris</i> Extract Inhibits the NF-κB pathway and Induces Apoptosis through MKK7-JNK Signaling Activation in TK-10 Human Renal Cell Carcinoma Soo Jung Park, Hyun-Jin Jang, In-Hu Hwang, Jung Min Kim, Eunbi Jo, Min-Goo Lee, Ik-Soon Jang and Jong Cheon Joo	465
<i>Aceriphyllum rossii</i> Exerts Lipid-Lowering Action in Both Normal and Hyperlipidemic Mice Mun Ock Kim, Jee Hee Seo, Eun Bin Kwon, Myung Ji Kang, Su Ui Lee, Dong Oh Moon, Mi Kyeong Lee, Chul-Ho Lee and Hyun Sun Lee	471

Continued inside backcover